



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\begin{aligned}
& + 2\log[\theta + \sqrt{(2 + \theta^2)}] \Big]_{\theta=0}^{\theta=60} = 3\sec\alpha \left\{ \frac{1}{2} [60\pi\sqrt{(2 + 3600\pi^2)} \right. \\
& \quad \left. + 2\log[60\pi + \sqrt{(2 + 3600\pi^2)}] - \frac{1}{2} [0\sqrt{2} + 2\log(0 + \sqrt{2})] \right\} \\
& = 3\sec\alpha \{ 30\pi\sqrt{(2 + 3600\pi^2)} + \log[60\pi + \sqrt{(2 + 3600\pi^2)}] - \log\sqrt{2} \} = 4540 \text{ feet.}
\end{aligned}$$

# DIOPHANTINE ANALYSIS.

116. Proposed by HARRY S. VANDIVER, Bala, Pa.

If  $n$  is an odd positive integer, and  $1, n, n', n'', \dots$  denote all its distinct divisors, then  $2^n > 2[n+1][n'+1][n''+1]\dots$

Solution by L. E. DICKSON, Ph. D., Assistant Professor of Mathematics, The University of Chicago.

There is a single exception  $n=3$ , for which  $2^3=2[3+1]$ . The corrected theorem may be proved by induction, using the following lemma:

If  $p$  is an odd prime number and  $d$  and  $\pi$  positive integers,

$$[1+d]^{p^\pi} > [1+d][1+pd][1+p^2d]\dots[1+p^\pi d],$$

except for  $d=1, \pi=1, p=3$ , the equality sign then holding.

For proof we apply  $p^\pi - 1 = p - 1 + p[p-1] + p^2[p-1] + \dots + p^{\pi-1}[p-1]$ .

$$\therefore [1+d]^{p^\pi} = [1+d][1+d]^{p-1}[1+d]^{p(p-1)}\dots[1+d]^{p^{\pi-1}(p-1)}.$$

But  $[1+d]^{t(p-1)} \geq [1+td]^{p-1} \geq 1 + [p-1]td + [td]^{p-1} \geq 1 + [p-1]td + td$ , if  $p > 2$ . The equality signs hold simultaneously only when  $t=d=1, p=3$ . Hence, for  $v > 2$ ,  $[1+d]^{t(p-1)} > 1 + ptd$  unless  $t=d=1, p=3$ , so that the lemma follows.

To prove the theorem by induction, we note that it is true for  $n=p^\pi > 3$ , in view of the lemma for  $d=1$ . Assume that it has been verified for  $n=p_1^{\pi_1}p_2^{\pi_2}\dots$ . We proceed to prove it true for  $N=np^\pi$ ,  $p$  being prime to  $n$ . We have

$$2^N = [2^n]^{p^\pi} \geq [(1+1)(1+n)(1+n)\dots]^{p^\pi}$$

$$\geq [(1+1)(1+p)\dots(1+p^\pi)][(1+n)(1+pn)(1+p^2n)\dots][(1+n')(1+p'n')\dots],$$

in view of the lemma. But the distinct divisors of  $N$  are

$$1, n, n', \dots, p, pn, pn', \dots, p^2, p^2n, \dots, p^\pi, p^\pi n, \dots$$

The theorem is therefore true for  $N$ .

118. Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Find the two least integral numbers such that their sum shall be a square and the sum of their squares a biquadrate.

Solution by G. B. M. ZERE, A. M., Ph. D., Parsons, W. Va.

Let  $x$  and  $y$  be the numbers, then for  $x+y=1, x^2+y^2=13^4$ ,  
 $x=120, y=-119$ .